

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE SEP 2008		2. REPORT TYPE		3. DATES COVERED 00-00-2008 to 00-00-2008	
4. TITLE AND SUBTITLE The International Arctic Buoy Programme (IABP): A Cornerstone of the Arctic Observing Network				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Washington,Polar Science Center,Applied Physics Laboratory,Seattle,WA,98105-6698				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES See also ADM002176. Presented at the MTS/IEEE Oceans 2008 Conference and Exhibition held in Quebec City, Canada on 15-18 September 2008.					
14. ABSTRACT see report					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 3	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

The International Arctic Buoy Programme (IABP): A Cornerstone of the Arctic Observing Network

Ignatius G. Rigor

Polar Science Center, Applied Physics Laboratory, University of Washington
Seattle, Washington, USA

Pablo Clemente-Colón

National/Naval Ice Center, Washington, D.C., USA

Ed Hudson

Meteorological Service of Canada, Edmonton, Alberta, Canada

Abstract—The Arctic has undergone dramatic changes in weather, climate and environment. It should be noted that many of these changes were first observed and studied using data from the IABP (<http://iabp.apl.washington.edu>). For example, IABP data were fundamental to Walsh et al. (1996) showing that atmospheric pressure has decreased (Figure 1), Rigor et al. (2000) showing that air temperatures have increased (Figure 2), and to Proshutinsky and Johnson (1997); Steele and Boyd, (1998); Kwok, (2000); and Rigor et al. (2002) showing that the clockwise circulation of sea ice and the ocean has weakened (Figure 1). All these results relied heavily on IABP data.

In addition to supporting these studies of climate change, the IABP observations are also used to validate satellite retrievals of environmental variables, to force, validate and initialize numerical models, and to forecast weather (e.g. Figure 3) and ice conditions. Over 600 papers have been written using data from the IABP. The observations and datasets of the IABP are one of the cornerstones for environmental forecasting and research in the Arctic.

Simply maintaining the network may be the biggest challenge for the IABP given the changes in climate. The winds tends to blow the buoys away from the Eurasian coast more quickly (Figure 1), and the decline of sea ice requires the development of more robust equipment that may survive the annual freeze/thaw cycles (Figure 4).

This talk summarizes the operations, some recent research, and the challenges facing IABP up through the International Polar Year, and beyond.




Figure 1. Mean field of SLP and ice motion for 1979–1986 (left), 1987–1994 (middle), and the difference between these two 8-year periods (right). These figures show a cyclonic anomaly in circulation (Adapted from Walsh, et al., 1996; Rigor, et al., 2002).




Figure 2. Seasonal Trends in surface air temperature derived from buoy data (Adapted from Rigor et al., 2000).




Figure 3. Analyzed field of surface air pressure for March 18, 2008.




Figure 4. Schematic of a prototype Airborne eXpendable Ice Buoy (AXIB), that designed to be deployed by air, and survive the annual freeze/thaw cycles in the larger areas of season ice in the Arctic Ocean. E.g. the map on the left shows the ice concentrations on October 12, 2007 just after the record minimum ice extent set in September 2007. Note the dearth of buoys in the areas of open, ice-free ocean (red arrow).